




Assessment of Diastolic Function

Challenging, but Can be Simple
ASE Echo Board Review Course
May 8th, 2018



Jae K. Oh, MD
Samsung Professor of CV Diseases

©2018 MPIMER | 3712003-1



Learning Objectives for Diastology After this talk, you will be able to

- Understand physiology and hemodynamics of diastole
- Know correlation between Echo diastolic parameters and underlying hemodynamics
- Classify and grade diastolic function
- Estimate filling pressure reliably in most patients at rest and with exercise
- Understand pitfalls of Echo diastolic function assessment



©2018 MPIMER | 3712003-2

Diastolic Filling with Relaxation



A Clinical Study of Left Ventricular Relaxation

YUZO HIROTA, M.D.

Circulation 1980


SUMMARY Left ventricular (LV) relaxation was studied in patients with hypertrophic cardiomyopathy (HCM, $n = 18$), congestive cardiomyopathy (CCM, $n = 11$), hypertensive heart disease (HHD, $n = 8$), coronary artery disease (CAD) without left ventricular (LV) asynergy ($n = 9$) and with LV asynergy ($n = 17$), mitral stenosis (MS, $n = 16$), and mitral regurgitation (MR, $n = 8$). The time constant T and peak negative dP/dt were used as indexes of LV relaxation, and 18 normal subjects served as controls.

The time constant T was higher in elderly patients among normal controls ($r = 0.652$, $p < 0.01$), which suggests that prolongation of relaxation is a phenomenon of aging. The normal value of the time constant T was 33 ± 8 msec (mean \pm SD), and that of peak negative dP/dt was 1864 ± 390 mm Hg/sec. The time constant T was significantly higher in HCM (64 ± 20 msec), CCM (56 ± 14 msec), CAD without asynergy (53 ± 16 msec), CAD with asynergy (57 ± 13 msec) and MS (47 ± 12 msec). Peak negative dP/dt was significantly lower in HCM (998 ± 303 mm Hg/sec), CCM (1060 ± 334 mm Hg/sec), CAD with asynergy (1370 ± 299 mm Hg/sec), MS (1367 ± 313 mm Hg/sec) and MR (1139 ± 305 mm Hg/sec).

Myocardial relaxation is one of the earliest manifestations of mechanical dysfunction of the human LV. The time constant tau (T) is higher in the elderly and patients with HCM, CAD, and cardiomyopathies.

Echo evaluation of diastolic function



Trans-mitral inflow velocity



©2018 MPIMER | 3712003-6

Diastolic Function Grading


Mitral Inflow (U curve)



Concept from Appleton and Hatle, 1985

©2018 MPIMER | 3712003-6

Mitral Inflow and Pulmonary Vein Flow Diastolic Function Assessment



Grade 1


Grade 2

Grade 3



©2018 MPIMER | 3712003-7


LV Relaxation by Cath and Echo τ (tau) vs e' (mitral annulus velocity)



Firstenberg et al: J Appl Physiol 90:299, 2001, Nagueh et al: JACC 1997
 Oki et al: AJC 1997, Sohn et al: JACC 1997, Ommen et al: Circ 2000 Opdahl et al: Circulation 119:2578, 2009, and more

©2018 MPIMER | 3712003-8


Assessment of LV Relaxation by Echo e' velocity reflects LV relaxation



Myocardial Relaxation is the Key for Diastole

©2018 MFMER | 3712003-9


Myocardial Relaxation (e')



©2018 MFMER | 3712003-10

Evaluation of Diastolic Function

Mitral Inflow and Annulus Velocity



Sohn et al: JACC, 1997



CP1204003-00
©2018 MFMER | 3712003-11


As LV filling pressure ↑

Mitral E ↑

Annulus E' ↓


E/E' ↑

Nagueh et al: JACC, 1997
Ommen et al: Circ, 2000



CP1204003-01
©2018 MFMER | 3712003-12

Estimation of LV Filling Pressures E/e' (Medial MV annulus)



Ommen SR et al: Circulation 102:1788, 2000




©2018 MFMER | 3712003-14

What are normal values for e' and E/e' ?



©2018 MFMER | 3712003-14

Echocardiographic reference ranges for normal cardiac Doppler data: results from the NORRE Study




Luis Caballero et al. EHJ 2015

©2018 MFMER | 3712003-16



ORIGINAL RESEARCH

Serial Doppler Echocardiography and Tissue Doppler Imaging in the Detection of Elevated Directly Measured Left Atrial Pressure in Ambulant Subjects With Chronic Heart Failure



Ritzema et al JACC Imaging 2011

©2018 MFMER | 3712003-16



ASE/EACVI GUIDELINES AND STANDARDS

Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging

Sherif F. Nagueh, Chair, MD, FASE,¹ Otto A. Smiseth, Co-Chair, MD, PhD,² Christopher P. Appleton, MD,¹
Benjamin F. Byrd, III, MD, FASE,¹ Hisham Dokainish, MD, FASE,¹ Thor Edvardsen, MD, PhD,²
Frank A. Flachskampf, MD, PhD, FESC,² Thierry C. Gillebert, MD, PhD, FESC,² Allan L. Klein, MD, FASE,¹
Patrizio Lancellotti, MD, PhD, FESC,² Paolo Marino, MD, FESC,² Jae K. Oh, MD,¹
Bogdan Alexandru Popescu, MD, PhD, FESC, FASE,² and Alan D. Waggoner, MHS, RDCS¹, *Houston, Texas;*

Four Major Diagnostic Parameters Normal Values


1. E' velocity ≥ 7 (med), 10 (lat) cm/s
2. E/e' ≤ 14 (Av), 15(Med)
3. TR velocity ≤ 2.8 m/sec
4. LAVI ≤ 34 mL/m²



JASE and EJ CV Imaging April 2016

©2016 MFMER | 3712003-16



New Criteria for Diastolic Function Assessment



Criteria for diagnosis of LV diastolic dysfunction in patients with normal LVEF in JASE 2016



©2016 MFMER | 3712003-20

Normal Diastolic Function




©2018 MFMER | 3712003-21

True Normal Diastolic Function




©2018 MFMER | 3712003-22

71 year old woman with LAVI = 39 mL/m²



71 year old woman with LAVI = 39 mL/m²




Reasons for LA enlargement

- Diastolic dysfunction
- Increased filling pressure
- Increased volume
- Athlete's heart
- Measurement error



©2018 MFMER | 3712003-26

Normal or Grade 1 Diastolic Dysfunction ? LAVI 28 mL/m²




E/e' NL < 14
e' ABNL < 7
LAVI Normal
TR NL < 2.8

©2018 MFMER | 3712003-26

New Criteria for Diastolic Function Assessment

In pts with normal LVEF $\geq 50\%$

- 1 – Septal e' velocity ≥ 7 cm/s or lateral e' velocity ≥ 10 cm/s
- 2 – Average $E/e' \leq 14$, 15 (Med)
- 3 – TR velocity ≤ 2.8 m/s
- 4 – LA volume index ≤ 34 mL/m²



≥ 3 Normal

2 and 2

≥ 3 Abnormal

Normal diastolic function

Indeterminate

Diastolic dysfunction

Normal Diastolic Filling Pressure

Criteria for diagnosis of LV diastolic dysfunction




Increased Diastolic Filling Pressure

LVEF



©2018 MFMER | 3712003-27


Diastolic Function?



e' ABNL < 7
 E/e' ABNL > 14
 LAVI ??
 TR NL > 2.8



©2018 MFMER | 3712003-28





Diastolic Function Assessment


Take Home Point #1

- LV myocardial relaxation is reduced in all stages of diastolic dysfunction
- Mitral annulus e' velocity reflects myocardial relaxation
- Normal e' = Normal diastolic function
- Algorithm #1 separates normal filling from elevated filling pressure
- Initial assessment of diastolic function is based on
 - E', E/e', TR velocity, and LAVI


Diastolic Function?





67 yo man with ischemic CM and HF
Gr. 1 dysfunction with normal filling pressure




67 yo man with ischemic CM and HF
Gr. 1 dysfunction with normal filling pressure






67 yo man with ischemic CM and HF Gr. 2 dysfunction with increased filling pressure



Ischemic Cardiomyopathy Echo Predictor STICH Trial (N=1511)



Lin et al. 2014 AHA

©2018 MFMER | 3712003-38

Distribution of 2D and Doppler Variables

		Elevated filling pressure (n=165)	Normal filling pressure (n=155)
Mitral E/A ratio ≤ 0.8 + E ≤ 50 cm/s		0	23
Mitral E/A ratio ≥ 2		53	5
None of the cutoff values met for the 3 variables in patients with diastolic dysfunction		15	70
3 abnormal	LAV >34 mL/m ² , E/e' >14 , and TRV >2.8 m/s	25	0
2 abnormal (2 of 3 listed)	LAV >34 mL/m ² , E/e' >14 , TRV <2.8 m/s	35	7
	LAV >34 mL/m ² , E/e' <14 , TRV >2.8 m/s	11	8
1 abnormal	LAV <34 mL/m ² , E/e' >14 , TRV >2.8 m/s	8	1
	LAV >34 mL/m ²	6	32
	E/e' >14	8	4
	TRV >2.8 m/s	4	5




Andersen et al: J Am Coll Cardiol 2017;69:1937-48


©2018 MFMER | 3712003-41

- Normal Filling Pressure
 - 3 normal parameters
 - LAVI >34 mL/m²
 - E/A ≤ 0.8 + E ≤ 50 cm/s
- Increased Filling pressure
 - E/A ≥ 2
 - E/e' > 14 + LAVI >34 mL/m²
 - 3 abnormal
 - 2 abnormal



23 YO with HCM





1. Grade 1
2. Grade 2
3. Grade 3
4. Possibly normal




©2018 MFMER | 3712003-42






Doppler Determination of LVEDP



EDP
(mm Hg)



Rossvoll and Hatle:
JACC, 1993

LVEDP can be increased with normal mean LV diastolic pressure

CP1057136-18
©2018 MFMER | 3712003-47

Diastolic Function Assessment Take Home Message #2

- Grade 1 diastolic dysfunction is the best pattern for the patients with Heart Failure
- Evidence for diastolic dysfunction needs an objective evidence
 - Hypertension
 - Hypertrophic CM
 - Old age

The best evidence is reduced relaxation
Reduced e'
L wave
Prolonged IVRT

Difficult Situations


- Assess diastolic function or filling pressure in
 - 2 normal and 2 abnormal
 - HCM
 - LBBB
 - MAC
 - Atrial Fibrillation
- Additional supportive parameters
 - Pulmonary vein
 - Valsalva
 - IVRT and timing intervals
 - Strain



©2018 MFMER | 3712003-48

67 year old woman with HPT and SOB

LAVI = 54 and TR= 2.8 m/s




1. $E' \pm$
2. E/e' NL
3. TR ≤ 2.8
4. LAVI ++



©2018 MFMER | 3712003-50





Indeterminate ?



©2018 MFMER | 3712003-81


Pulmonary Vein Velocity

PVs decreases as filling pressure increases



©2018 MFMER | 3712003-82

Valsalva Maneuver : E/A reduced > 0.5 Grade 2 Dysfunction




Evaluation of Left Ventricular Filling Pressures by Doppler Echocardiography in Patients With Hypertrophic Cardiomyopathy Correlation With Direct Left Atrial Pressure Measurement at Cardiac Catheterization

Jeffrey B. Geske, MD; Paul Sorajja, MD; Rick A. Nishimura, MD; Steve R. Ommen, MD

Conclusions—In 100 symptomatic patients with HCM, Doppler echo estimates of LV filling pressure correlate modestly with direct measurement of LAP. Given the complex nature of diastolic dysfunction in HCM, precise characterization of LV filling pressure in an individual patient cannot be determined with the use of these noninvasive parameters. (*Circulation*. 2007;116:2702-2708.)



Medial E/e' Ratio Versus Mean LAP



Geske et al: Circulation; 116:2702, 2007

©2018 MFMER | 3712003-95

Mean LAP vs Medial E-e' ratio Hypertrophic CM



Geske et al: Circulation; 116:2702, 2007


©2018 MFMER | 3712003-96

Diastolic Function Evaluation in HCM

- E' velocity is reduced in almost all patients
- E/e' predicts clinical outcome
- Use following parameters (ASE 2016 Guideline)
 - $E/e' > 15$
 - LAVI > 34 mL/m²
 - TR velocity > 2.8 m/sec
 - PV Ar-A duration ≥ 30 msec
- The majority rules





©2018 MEDICAL J. 3712003-97



72 yo woman with HCM




72 yo woman with apical HCM Grade 2 dysfunction with LAVI 37 mL/m²




Mitral annulus e' velocity

- ASE/EACVI recommends average value
- E' from one location is acceptable
- We need a caution in using e'
 - Primary pulmonary hypertension
 - Pacemaker
 - LBBB
 - Wall motion abnormality
 - Mitral annulus calcification
 - Hypertrophic CM


67 yo woman with LBBB




67 yo with LBBB



Mitral annulus calcification and TAVR



Mitral annulus calcification and TAVR



Mitral annulus e' velocity vs MAC

Mean age 73 years

Variable	Group 1 n=79 no MAC	Group 2 n=38 mild MAC	Group 3 n=38 mod-severe MAC	P for trend
Agatston Score	0	1-119	>119	
Septal e'	5.96±1.82	5.15±1.56	5.05±1.93	0.01
Lateral e'	7.37±2.44	6.89±2.71	6.28±1.81	0.01
Average e'	6.63±2	6.02±1.79	5.67±1.69	0.01
E/avg e' ratio	13±4.93	15±8.95	18±8.26	<0.001

LV diastolic parameters are altered in the presence of MAC. This could be due to direct effects of MAC or might reflect truly reduced diastolic function. Interpretation of diastolic parameters in patients with MAC should be performed with caution.



Codolosa et al: Am J Cardiol 2016;117:847-852

©2018 MFMER | 3712003-68

JACC: CARDIOVASCULAR IMAGING
© 2017 BY THE AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION
PUBLISHED BY ELSEVIER

VOL. ■, NO. ■, 2017
ISSN 1936-878X/\$36.00
<http://dx.doi.org/10.1016/j.jcmg.2016.10.017>

Doppler Echocardiography for the Estimation of LV Filling Pressure in Patients With Mitral Annular Calcification

Muaz M. Abudiab, MD, Lakshmi H. Chebrolu, MD, Robert C. Schutt, MD, Sherif F. Nagueh, MD, William A. Zoghbi, MD


Representative Images of Patients With Mitral Annular Calcification



Mild



Moderate




Severe



Abudiab et al: Am Coll Cardiol Img, 2017

©2018 MFMER | 3712003-68

Correlation of Selected Doppler Variables With Left Ventricular Filling Pressure



Abudiab et al: Am Coll Cardiol Img, 2017

©2018 MFMER | 3712003-67


Proposed Clinical Algorithm for Estimation of Left Ventricular Filling Pressure in Subjects With Mitral Annular Calcification

Initial Cohort (n=50):

Sensitivity: 81%
Specificity: 100%
PPV: 100%
NPV: 67%

Total Cohort (n=71):

Sensitivity: 85%
Specificity: 95%
PPV: 97%
NPV: 78%



Abudiab et al: Am Coll Cardiol Img, 2017

©2018 MFMER | 3712003-68

Diastolic Function in A. Fib




- DT < 160 msec (with reduced EF)
- DT < 130 msec poor survival (Hurley, Oh)
- Other measurements
 - E acceleration > 1900 cm/sec²
 - IVRT ≤ 65 msec
 - E/e' ≥ 11
 - IVRT/ T E-e'
 - TR velocity



©2018 MFMER | 3712003-66

Mitral Annulus Velocity in the Evaluation of Left Ventricular Diastolic Function in Atrial Fibrillation

Dae-Won Sohn, MD, Jong-Min Song, MD, Joo-Hee Zo, MD, In-Ho Chai, MD, Hyo-Soo Kim, MD, Hong-Gu Chun, MA, and Hee-Chan Kim, PhD, *Seoul, Korea*




JASE 1999



©2018 MFMER | 3712003-70

Atrial Fibrillation


Variation in E velocities : NL Pressure



©2018 MFMER | 3712003-71


Atrial Fibrillation

Variation in E velocities and $E/e' < 11$




©2018 MFMER | 3712003-72

66 yo woman with dyspnea and EF 20%




66 yo woman with HF and severe MR LVEF 20%



Doppler Estimation of Left Ventricular Filling Pressure in Sinus Tachycardia

A New Application of Tissue Doppler Imaging

Sherif F. Nagueh, MD; Issam Mikati, MD; Helen A. Kopelen, RDMS; Katherine J. Middleton, RCT; Miguel A. Quiñones, MD; William A. Zoghbi, MD



Nagueh et al Circulation 1998

©2018 MFMER | 3/12/2003-76

My Recommendations

- For patients with reduced EF (<35%) or with preserved EF with **known diastolic dysfunction**, evaluate diastolic function based on E/A ratio
- For all other patients, based on the 4 parameters
 - Normal : ≥ 3 normal (for patient's age)
 - Abnormal : ≥ 3 abnormal (grade 2 or 3 based on E/A)
 - Indeterminate : Need help from PV, IVRT, Valsalva, Time interval, Exercise, Strain Imaging




©2018 MFMER | 3/12/2003-76

My recommendation based on e' velocity LV relaxation is the key for normal diastole

- True Normal
 - Medial e' >10 cm/sec or Lateral e' > 15 cm/sec
- Age-related Normal
 - Medial e' 7-10 cm/s or Lateral e' 10-15 with normal TR
- Abnormal
 - Medial e' < 7 cm/s or lateral e' <10 cm/s
 - Grade 1, 2, and 3 based on E/e', TR, and LAVI



©2018 MFMER | 3712003-27



©2018 MFMER | 3712003-78

Thank you!
Oh.jae@mayo.edu